



# Trenolin® Super<sup>PLUS</sup>

Highly active, depectidase-free, liquid pectinase


## Product description

Trenolin® Super<sup>PLUS</sup> is a liquid formulation containing various active pectinases for rapid degradation of the complex pectin molecule. The enzyme is used in mashes, juice, young wine and Süßreserve. Trenolin® Super<sup>PLUS</sup> is depectidase-free.

The benefits of Trenolin® Super<sup>PLUS</sup> are emphasized by the optimised recipe and increased activity. These are:

- Increased free run juice
- Faster and more compact deposition of sediments
- Improved filtration and clarification

Other benefits	<ul style="list-style-type: none"> <li>• Increased press capacity</li> <li>• Reduced press and processing times</li> <li>• Reduction of tannin discharge and subsequent treatments</li> <li>• Preventative improvement of filtration performance in wine</li> </ul>
Recommended for	<ul style="list-style-type: none"> <li>• All types of white, rosé and red wine</li> </ul>



In accordance with Regulation (EC) no. 934/2019. Laboratory tested for purity and quality.

**PLUS** - new formula means additional benefit!

The following properties are characteristic of the Trenolin®<sup>PLUS</sup> series of enzymes:

- Recipe optimised for modern wine-making
- Depectidase-free (free from cinnamyl esterase activity)
- No preservatives

## Dosage

	mL/100 L or 100 kg mash		
Grapes	Approx. 7 - 15	Young wine	Approx. 3 - 5
Mash	Approx. 5 - 10	Süßreserve	Approx. 5 - 10
Juice	Approx. 5		
Temperature	The recommended doses are based on a temperature of 15 - 17 °C.		
Application time	Usually at least 1 hour depending on the variety's pectin content and framework conditions.		
Tip	Effectiveness depends on the quantity of product, temperature and reaction time. The treatment temperature should be more than 12 °C and preferably more than 15 °C. The higher the temperature, the more effective the enzyme. The natural upper limit is 55 °C.		
Application	Dilute the appropriate quantity of enzyme per tank with a little liquid to achieve better and more even distribution.		
Attention!	Bentonites deactivate the enzyme, so bentonite should only be used after the pectin has been fully degraded (pectin test).		

## Storage

Store in a cool place. Packs which have been opened should be immediately tightly sealed and used up as soon as possible.

ERBSLÖH Geisenheim GmbH • Erbslöhstraße 1 • 65366 Geisenheim, Germany  
Tel.: +49 6722 708-0 • Fax: +49 6722 6098 • info@erbsloeh.com • www.erbsloeh.com

**ERBSLÖH**  
Fortschritt macht Zukunft®

The application recommendations given herein describe the intended use of the product as a processing aid or additive as part of a good manufacturing practice. Only this application can lead to a food safety of the final product. However, please note: Our technical product leaflets are based on our current knowledge and experience. They have to be seen as general information on our products only. Due to the imponderabilities of treating natural products and the potential prior treatment we cannot accept any liability. Accordance with all national laws and regulations for use of our products has to be ensured by each user. All data is therefore provided without any warranty. All information is subject to change without prior notice. Our general terms of business apply, please refer to www.erbsloeh.com.  
Version 004-11/2019 EH – printed 28.11.2019